404000mE

408000mE

410000mE

412000mE

404000mE

406000mE

408000mE

410000mE

412000mE



LAND RESOURCES of the **UNDOOLYA ROCKY HILL** AGRICULTURAL BLOCK

For further information contact:

Department of Environment, Parks and Water Security Director, Land Assessment, Rangelands Division

Ph. (08) 8999 4478 Email: rangelands@nt.gov.au Web: http://denr.nt.gov.au Level 3, Goyder Centre, 25 Chung Wah Terrace, Palmerston, Northern Territory of Australia.

> NR Maps: https://nrmaps.nt.gov.au Map Reference: DENR2020085 Undoolya-Rocky-Hill Land-Resources Map

LAND UNIT DESCRIPTIONS

Small gravelly mesa and sideslopes. Gravelly structured soils (Red Dermosols). Mixed species sparse tussock grassland with forb species.

- Bare areas among sandy rises and gravelly pediments. Undescribed. Largely bare or devoid of vegetation. Occasional pockets of Senna sp. open shrubland.
- Undulating to rolling rises. Red sandy soils (Red-Orthic Tenosols). Acacia aneura with Acacia estrophiolata Mid high open woodland.

LOW RISES

RISES

- Drainage depressions within sandy rises, low rises and undulating plains. Red loamy soils (Red Kandosols). Acacia aneura and Acacia estrophiolata Mid high open woodland. Drainage depressions within sandy rises, low rises and undulating plains. Red loamy soils (Red Kandosols). Acacia estrophiolata with Acacia aneura and Eucalyptus coolabah
- Low rises and gently undulating plains. Red loamy soils (Red Kandosols). Acacia estrophiolata with Hakea divaricata and Corymbia apparrerinja Low open woodland.

subsp. arida Tall open woodland.

Dermosols). Vegetation has not been described.

PLAINS

- Mainly level floodplains with some raised areas of sandy and loamy deposits and some small depressions. Red earths (Red Kandosols and Dermosols). A number of various communities and bare areas exist and a dominant structure cannot be determined. Dissected margins of the flooplain; areas with moderate to large gullies. Red earths
- FP-DM (Red Kandosols and Dermosols). Vegetation has not been described. Floodplain with surface erosion and minor gullies. Red earths (Red Kandosols and
- Flat area of floodplain beside sandy rises overlying rock. Red brown loamy soils (Lithic FP-Sal Calcic Calcarosols). Largely bare or devoid of vegetation.
- Floodplains; minor areas of narrow elongated sandy deposits up to 50m wide and 1.2m FPa high. Red brown loamy sands (Red-Orthic Tenosols). Acacia estrophiolata Tall open

Floodplains; minor areas of narrow elongated sandy deposits up to 50m wide and 1.2m

- FPa/FP high. Red brown loamy sands (Red-Orthic Tenosols). Acacia estrophiolata Tall open Bare areas of the floodplain; often slightly lower lying areas; can occur as very shallow basins and as closed depressions. Red earths (Red Dermosols and Kandosols). Largely
- bare or devoid of vegetation. Drainage lines and low lying areas. Red earths (Red Dermosols). Eucalyptus coolabah subsp. arida with Acacia aneura and Acacia estrophiolata Mid high open woodland.
- Undulating very slightly raised irregular sandy area of floodplain. Red sandy loams (Orthic Tenosols). Acacia georginae Mid high woodland. Floodplain areas adjacent to drainage areas. Red earths (Red Kandosols and Dermosols). Eucalyptus coolabah subsp. arida Dwarf open woodland (largely E. coolabah seedlings).
- Slightly elevated floodplain. Red loamy soils (Red Dermosols). Eucalyptus coolabah subsp. arida with Acacia estrophiolata and Acacia tetragonophylla Low open woodland/Tall sparse shrubland.
- Small areas on floodplain that are very slightly raised or have a soft surface. Red earths (Red Kandosols). Acacia estrophiolata with Acacia kempeana and Hakea divaricata Tall
- Small areas on floodplain that are very slightly raised or have a soft surface. Red earths FPs/FP (Red Kandosols). Acacia estrophiolata with Acacia kempeana and Hakea divaricata
- Lower lying areas of floodplain. Red earths (Red Kandosols). Acacia estrophiolata with Eucalyptus coolabah subsp. arida and Hakea divaricata Mid high open woodland.
- Gravelly pediment. Gravelly soils (Calcic Calcarosols). Largely bare or devoid of GP-Sal Low lying sections of pediment; downslope from gravelly rises and salinised gravelly
- pediment; near low lying areas of alluvial plain. Red sandy or loamy soils (Red Kandosols) Senna artemisioides subsp. filifolia with Acacia aneura and Hakea leucoptera Tall open Mid to upper sections of pediment below gravelly rises; rock exposed in some areas.
- Red loamy soils (Red Kandosols). Eremophila freelingii, Eremophila duttonii and Senna artemisioides subsp. alicia Tall open shrubland. Relict prior streams and floodout deposits; elevated up to 2m above adjacent floodplains.
 - Sandy soils (Red-Orthic Tenosols). Acacia estrophiolata Mid high open woodland.

ALLUVIAL PLAINS Lower lying areas of alluvial plains. Red loamy soils (Red Kandosols). Acacia aneura with

- Acacia estrophiolata Mid high woodland.
 - Level alluvial plains. Red loamy soils (Red Kandosols). Acacia aneura with
 - Acacia estrophiolata Tall woodland.

DRAINAGE Stream channels; banks and stream bed, some channel benches. Sandy to loamy soils SC

(Tenosols). Eucalyptus camaldulensis and E. coolabah subsp. arida Open woodland on larger banks and channel benches. Some areas of Eucalyptus coolabah open woodland with a grass understorey, other areas with sparse shrubs on banks.

Example of Land Unit Descriptions - Landform description - Landform — Soil description PLAINS Mainly level floodplains with some raised areas of sandy and loamy deposits and some small depressions. Red earths (Red Kandosols and Dermosols). A number of various communities and bare areas exist and a dominant structure cannot be determined. - Land unit Vegetation description

MAP DISCLAIMER:

Land resource information has been derived from aerial photograph interpretation and field data describing landform, soil and vegetation. Mapping has been collected according to the national standards and prepared at a scale of 1:25 000. Enlarging this map beyond this scale will not provide further detail.

A site inspection should always accompany mapping for specific areas.

BIBLIOGRAPHIC REFERENCE:

Tucker J. Richard, Whitehouse M. Diana and King J. Daryl 1999. Land Resources of The Undoolya Rocky Hill Agricultural Block. A Reconnaissance Land Resource Survey to Assess Potential areas for Horticultural Development.

Natural Resources Division, Department of Lands, Planning and Environment, Alice Springs, NT.

TECHNICAL REFERENCES:

McDonald, R.C., Isbell, R.F., Speight, J.G., Walker, J. and Hopkins, M.S. (1990). Australian Soil and Land Survey Field Handbook. 2nd Edition. Inkata Press, Melbourne.

> Isbell R.F. (1996). The Australian Soil Classification. CSIRO Publishing, Melbourne.